

gradually decreased over a period of approximately two weeks towards an OR of one (2).

The possible contribution of airway infections, which occur more frequently in cold winter months than in summer, have not been listed by Spencer et al. (1) among the possible explanations for the higher incidence of AMIs in winter time. Evidence from smaller hospital-based studies has already previously indicated that an association between acute chest infections and AMI might exist (3-5). The pathophysiologic mechanism of such an association is speculative and could involve changes in circulating clotting factors, an increased risk for acute rupture of arteriosclerotic plaque during a chest infection, or a variety of other mechanisms. The role of acute and chronic infections in the etiology of coronary heart disease and AMI needs to be better understood; it might be one of the keys towards an explanation why more AMIs occur in winter than in summer.

CHRISTOPH R. MEIER, PhD, MSc
*Boston Collaborative Drug Surveillance Program
Boston University School of Medicine
11 Muzzey Street
Lexington, Massachusetts 02421*

References

1. Spencer FA, Goldberg RJ, Becker RC, Gore JM. Seasonal Distribution of Acute Myocardial Infarction in the Second National Registry of Myocardial Infarction. *J Am Coll Cardiol* 1998;31:1226-33.
2. Meier CR, Jick SS, Derby LE, Vasilakis C, Jick H. Acute respiratory-tract infections and risk of first-time myocardial infarction. *Lancet* 1998;351:1467-71.
3. Pesonen E, Siitonen O. Acute myocardial infarction precipitated by infectious disease. *Am Heart J* 1981;101:512-3.
4. Spodick DH, Flessas AP, Johnson MM. Association of acute respiratory symptoms with onset of acute myocardial infarction: prospective investigation of 150 consecutive patients and matched control patients. *Am J Cardiol* 1984;53:481-2.
5. Mattila KJ. Viral and bacterial infections in acute myocardial infarction. *J Intern Med* 1989;225:293-6.

Anticoagulation in Dilated Cardiomyopathy

We read with interest the recent review by Koniaris et al. (1) concerning anticoagulation in dilated cardiomyopathy. Evidence from published studies does not convincingly demonstrate the benefits of

anticoagulation in patients with dilated cardiomyopathy. As described by Koniaris et al., (1) a prospective, randomized clinical trial of long-term anticoagulation in patients with dilated cardiomyopathy is feasible. However, studies employing hemostatic markers indicating a biochemical imbalance between procoagulant and anticoagulant mechanisms in the blood may be useful to evaluate the appearance of thrombotic phenomena in these patients (2). By measuring plasma levels of hemostatic markers, we previously found that, in patients with dilated cardiomyopathy, plasma levels of fibrinopeptide A and thrombin-antithrombin III complex, markers of coagulation activation and thrombin generation, were significantly higher than those in normal subjects (3). Their levels showed a positive correlation with left ventricular end-diastolic volume and a negative correlation with fractional shortening of the left ventricle. Although Koniaris et al. (1) suggested that aspirin monotherapy may be beneficial for risk reduction of thromboembolism in patients with dilated cardiomyopathy, plasma levels of platelet factor-4 and beta-thromboglobulin, markers of platelet activation, were not elevated in these patients compared with normal subjects. In addition, we previously observed that aspirin monotherapy suppresses platelet function, but does not affect coagulation activity (4). These findings support the premise that anticoagulant therapy, rather than antiplatelet therapy, is more effective for the prevention of systemic embolism in patients with dilated cardiomyopathy, particularly in those with severe left ventricular enlargement and dysfunction.

UICHI IKEDA, MD
KEIJI YAMAMOTO, MD
KAZUYUKI SHIMADA, MD
*Department of Cardiology
Jichi Medical School
Minamikawachi-Machi
Tochigi 329-0498, Japan*

References

1. Koniaris LS, Goldhaber SZ. Anticoagulation in dilated cardiomyopathy. *J Am Coll Cardiol* 1998;31:745-8.
2. Ikeda U, Yamamoto K, Shimada K. Biochemical markers of coagulation activation in mitral stenosis, atrial fibrillation and cardiomyopathy. *Clin Cardiol* 1997;20:7-10.
3. Yamamoto K, Ikeda U, Furuhashi K, Irokawa M, Nakayama T, Shimada K. The coagulation system is activated in idiopathic cardiomyopathy. *J Am Coll Cardiol* 1995;25:1634-40.
4. Yamamoto K, Ikeda U, Fukazawa H, Shimada K. Effects of aspirin on status of thrombin generation in atrial fibrillation. *Am J Cardiol* 1996;77:528-30.